- Abdullah, M.M., Lee, S.Y., 2017. Structure of mangrove meiofaunal assemblages associated with local sediment conditions in subtropical eastern Australia. Estuarine, Coastal and Shelf Science 198, 438–449.
- Adão, H., Alves, A., Patrício, J., Neto, J.M., Costa, M.J., Marques, J.C., 2009. Spatial distribution of subtidal Nematoda communities along the salinity gradient in two Southern European estuaries (Portugal). Acta Oecologica 35, 287–300.
- Alkemade, R., Wielemaker, A., Hemminga, M.A., 1992. Stimulation of decomposition of *Spartina anglica* leaves by the bacterivorous marine nematode *Diplolaimelloides bruciei* (Monhysteridae). Journal of Experimental Marine Biology and Ecology 159(2), 267–278.
- Allison, M.A., Khan, S.R., Goodbred, S.L.Jr., Kuehl, S.A., 2003. Stratigraphic evolution of the late Holocene Ganges–Brahmaputra lower delta plain. Sedimentary Geology 155, 317–342.
- Allouche, M., Nasri, A., Harrath, A.H., Mansour, L., Beyrem, H., Boufahja, F., 2020. Migratory behavior of free-living marine nematodes surrounded by sediments experimentally contaminated by mixtures of polycyclic aromatic hydrocarbons. Journal of King Saud University-Science 32, 1339–1345.
- Alloy, M.M., Boube, I., Griffitt, R.J., Oris, J.T., Roberts, A.P., 2015. Photo-induced toxicity of Deepwater Horizon slick oil to blue crab (*Callinectes sapidus*) larvae. Environmental Toxicology and Chemistry 34, 2061–2066.
- Alongi, D.M., 1985. Effects of physical disturbance on population dynamics and tropic interactions among microbes and meiofauna. Journal of Marine Research 43, 351– 364.
- Alongi, D.M., 1987a. Intertidal zonation and seasonality of meiobenthos in tropical mangrove estuaries. Marine Biology 95, 447–458.
- Alongi, D.M., 1987b. The influence of mangrove derived tannins on intertidal meiobenthos in tropical estuaries. Oecologia 71, 537–540.

- Alongi, D.M., 1990. The ecology of tropical soft-bottom benthic ecosystem. Oceanography and Marine Biology: An Annual Review 28, 381–496.
- Alongi, D.M., 1998. Coastal Ecosystem Processes. New York, USA: CRC Press, 419 pp.
- Alongi, D.M., 2002. Present state and future of the world's mangrove forests. Environmental conservation 29, 331–349.
- Al-Sofyani, A., El-Sherbiny, M., 2018. Meiobenthic assemblage of the grey mangrove (*Avicennia marina*) along the Saudi Arabian coast of the Red Sea with emphasis on free-living nematodes. Oceanological and Hydrobiological Studies 47(4), 359–375.
- Alsterberg, C., Hulth, S., Sundback, K., 2011. Response of a shallow-water sediment system to warming. Limnology and Oceanography 56, 2147–2160.
- Altaff, K., Sugumaran, J., Naveed, Md. S., 2005. Impact of tsunami on meiofauna of Marina Beach, Chennai, India. Current Science 89(1), 34–38.
- Alves, A.S., Adão, H., Patrício, J., Neto, J.M., Costa, M.J., Marques, J.C., 2009. Spatial distribution of subtidal meiobenthos along estuarine gradients in two southern European estuaries (Portugal). Journal of the Marine Biological Association of the United Kingdom 89, 1529–1540.
- Alves, A.S., Adão, H., Ferrero, T.J., Marques, J.C., Costa, M.J., Patrício, J., 2013. Benthic meiofauna as indicator of ecological changes in estuarine ecosystems: the use of nematodes in ecological quality assessment. Ecological Indicators 24, 462–475.
- Alves, A.S., Verissimo, H., Costa, M.J., Marques, J.C., 2014. Taxonomic resolution and biological traits analysis (BTA) approaches in estuarine free-living nematodes. Estuarine, Coastal and Shelf Science 138, 69–78.
- Alves, A.S., Caetano, A., Costa, J.L., Costa, M.J., Marques, J.C., 2015. Estuarine intertidal meiofauna and nematode communities as indicator of ecosystem's recovery following mitigation measures. Ecological Indicators 54, 184–196.
- Anderson, R.S., De Henau, A.M., 1980. An assessment of the meiobenthos from nine mountain lakes in western Canada. Hydrobiologia 70, 257–264.

- Anderson, M.J., Gorley, R.N., Clarke, K.R., 2008. PERMANOVA A+ for PRIMER: Guide to Software and Statistical Methods. PRIMER-E, Plymouth, UK.
- Anila Kumary, K.S., 2008. Diversity of meiobenthic nematodes in the Poonthura estuary (southwest coast of India). Journal of the Marine Biological Association of India 50(1), 23–28.
- Annamalai, H., Sperber, K.R. 2016. South Asian Summer Monsoon Variability in a Changing Climate. In: de Carvalho, L., Jones, C., (Eds.), The Monsoons and Climate Change. Springer Climate, Springer, Cham.
- Annandale, N., 1907. The fauna of the Brackish Ponds at Port Canning, Lower Bengal, 1. Introduction and Preliminary account of the fauna. Records of the Indian Museum 1(1), 35–43.
- Annapurna, C., Bhanu, C.V., Rao, M.S., Sivalakshmi, M.V., Cooper, L.M.G., Rao, Y.K., 2012. Free-living nematodes along the continental slope off northeast coast of India. Journal of the Marine Biological Association of India 54(2), 52–60.
- Ansari, Z.A., Parulekar, A.H., 1981. Meiofauna of the Andaman Sea. Indian Journal of Marine Sciences 10(3), 285–288.
- Ansari, Z.A., Parulekar, A.H., 1993. Distribution, abundance and ecology of the meiofauna in a tropical estuary along the west coast of India. Hydrobiologia 262, 115–126.
- Ansari, Z.A., Sreepada, R.A., Matondkar, S.G.P., Parulekar, A.H., 1993. Meiofauna stratification in relation to microbial food in a tropical mangrove mudflat. Journal of Tropical Ecology 34, 63–75.
- Ansari, Z.A., Gauns, M.U., 1996. A quantitative analysis of fine scale distribution of intertidal meiofauna in response to food resources. Indian Journal of Marine Sciences 25(3), 259–263.
- Ansari, Z.A., Ingole, B.S., Parulekar, A.H., 1996. Benthos of the EEZ of the India. In: Qasim, S.Z., Roonwal, G.S., (Eds), India's Exclusive Economic Zone. Omega Scientific Publishes, New Delhi, pp. 74–86.

- Ansari, Z.A., Parulekar, A.H., 1998. Community structure of meiobenthos from a tropical estuary. Indian Journal of Marine Sciences 27, 362–366.
- Ansari, A., Rivonkar, C.U., Sangodkar, U.M., 2001. Population fluctuation and vertical distribution of meiofauna in a tropical mudflat at Mandovi estuary, west coast of India. Indian Journal of Geo-Marine Sciences 30, 237–245.
- Ansari, K.G.M.T., Lyla, P.S., Khan, S.A., Manokaran, S., Raja, S., 2013. Community structure of harpacticoid copepods from the southeast continental shelf of India. Proceedings of the International Academy of Ecology and Environmental Sciences 3(2), 87–100.
- Ansari, K.G.M.T., Lyla, P.S., Khan, S.A., Bhadury, P., 2015. Diversity patterns of freeliving marine nematodes in the southwest continental shelf off Bay of Bengal and their link to abiotic variables. Marine Ecology 37, 631–644.
- Arieli, R.N., Almogi-Labin, A., Abramovich, S., Herut, B., 2011. The effect of thermal pollution on benthic foraminiferal assemblages in the Mediterranean shoreface adjacent to Hadera power plant (Israel). Marine Pollution Bulletin 62, 1002–1012.
- Arlt, G., Muller, B., Warnack, K.H., 1982. On the distribution of meiofauna in the Baltic Sea. Hydrobiologia 73, 297–308.
- Armenteros, M., Martin, I., Williams, J.P., Creagh, B., Gonzalez-Sanson, G., Capetillo, N., 2006. Spatial and temporal variations of meiofaunal communities from the western sector of the Gulf of Batabano, Cuba. I. Mangrove system. Estuaries and Coasts 29, 124–132.
- Armenteros, M., Williams, J.P., Creagh, B., Capetillo, N., 2008a. Spatial and temporal variations of meiofaunal communities from the western sector of the Gulf of Batabanó, Cuba: III. Vertical distribution. Revista de Biologia Tropical 56(3), 1127– 1134.
- Armenteros, M., Perez-Garcia, J.A., Perez-Angulo, A., Williams, J.P., 2008b. Efficiency of extraction of meiofauna from sandy and muddy marine sediments. Revista de Investigaciones Marinas 29(2), 113–118.

- Armenteros, M., Ruiz-Abierno, A., Fernández-Garcés, R., Pérez-García, J.A., Díaz-Asencio, L., Vincx, M., Decraemer, W., 2009. Biodiversity patterns of free-living marine nematodes in a tropical bay: Cienfuegos, Caribbean Sea. Estuarine, Coastal and Shelf Science 85, 179–189.
- Arnberg, M., Calosi, P., Spicer, J.I., Taban, I.C., Bamber, S.D., Westerlund, S., Vingen, S., Baussant, T., Bechmann, R.K., Dupont, S., 2018. Effects of oil and global environmental drivers on two keystone marine invertebrates. Scientific Reports 8(1), 17380.
- Artois, T., Fontaneto, D., Hummon, W.D., McInnes, S.J., Todaro, M.A., Sørensen, M.V.,
  Zullini, A., 2011. Ubiquity of microscopic animals? Evidence from the morphological approach in species identification. In: Fontaneto, D., (Eds),
  Biogeography of Microscopic Organisms: Is Everything Small Everywhere?
  Cambridge University Press, Cambridge, pp. 244–283.
- Attri, S.D., Tyagi, A., 2010. Climate Profile of India; Met. Monograph No. Environment Meteorology – 01/2010. India Meteorological Department, Ministry of Earth Sciences, Government of India, 129p.
- Austen, M.C., Warwick, R.M., 1989a. Meiobenthic and macrobenthic community structure along a putative pollution gradient in South Portugal. Marine Pollution Bulletin 20, 398–404.
- Austen, M.C., Warwick, R.M., 1989b. Comparison of univariate and multivariate aspects of estuarine meiobenthic community structure. Estuarine, Coastal and Shelf Science 29, 23–42.
- Austen, M.C., McEvoy, A.J., Warwick, R.M., 1994. The specificity of meiobenthic community response to different pollutants: results from microcosm experiments. Marine Pollution Bulletin 28, 557–563.
- Austen, M.C., Lambshead, J., Hutchings, P., Boucher, G., Heip, C., King, G., Koikel, Smith, C.R., Snelgrove, P.V.R., 2002. Biodiversity links above and below the marine sediment-water interface that may influence community stability. Biodiversity and Conservation 11, 113–136.

- Baliarsingh, S.K., Srichandan, S., Sahu, K.C., Lotliker, A.A., 2015. First record of *Desmoscolex falcatus* (Nematoda: Adenophorea: Desmoscolecida: Desmoscolecidae) from Rushikulya estuary, Odisha, India. Indian Journal of Geo-Marine Sciences 44(4), 519–521.
- Balsamo, M., Albertelli, G., Ceccherelli, V.U., Coccioni, R., Colangelo, M.A., Curini-Galletti, M., Danovaro, R., 2010. Meiofauna of the Adriatic Sea: current state of knowledge and future perspectives. Chemical Ecology 26(1), 45–63.
- Balsamo, M., Semprucci, F., Frontalini, F., Coccioni, R., 2012. Meiofauna as a tool for marine ecosystem biomonitoring marine ecosystems. In: Antonio, C., (Eds.), Marine ecosystems. In Tech, Rijeka, pp. 77–104.
- Barranguet, C., 1997. The role of microphyto benthic primary production in a Mediterranean mussel culture area. Estuarine, Coastal and Shelf Science 44, 753–765.
- Bauer, J.E., Cai, W.J., Raymond, P.A., Bianchi, T.S., Hopkinson, C.S., Regnier, P.A.G., 2013. The changing carbon cycle of the coastal ocean, Nature 504, 61–70.
- Beck, M.W., Heck, K., Able, K., Childers, D., Egglestone, D., Gillanders, B., Halpern, B., Hays, C., Hoshino, K., Minello, T., Orth, R., Sheridan, P., Weintein, M., 2001. The identification, conservation and management of estuarine and marine nurseries for fish and invertebrates. Bioscience 51, 633–641.
- Behrenfeld, M.J., O'Malley, R.T., Siegel, D.A., McClain, C.R., Sarmiento, J.L., Feldman, G.C., Milligan, A.J., Falkowski, P.G., Letelier, R.M., Boss, E.S., 2006. Climatedriven trends in contemporary ocean productivity. Nature 444(7120), 752–755.
- Bell, S.S., 1980. Meiofauna-macrofauna interactions in a high salt marsh habitat. Ecological monographs 50, 487–505.
- Bernardello, M., Secco, T., Pellizzato, F., 2006. The changing state of contamination in the lagoon of Venice. Part 2: heavy metals. Chemosphere 64, 1334–1345.
- Beyrem, H., Mahmoudi, E., Essid, N., Hedfi, A., Boufahja, F., Aissa, P., 2007. Individual and combined effects of cadmium and diesel on a nematode community in a laboratory microcosm experiment. Ecotoxicology and Environmental Safety 68, 412–418.

- Bezerra, T.N., Decraemer, W., Eisendle-Flöckner, U., Hodda, M., Holovachov, O., Leduc, D., Miljutin, D., Mokievsky, V., Peña Santiago, R., Sharma, J., Smol, N., Tchesunov, A., Venekey, V., Zhao, Z., Vanreusel, A., 2020. Nemys: World Database of Nematodes. Accessed at http://nemys.ugent.be on 2020-04-26. doi:10.14284/366.
- Bhadury, P., Austen, M.C., Bilton, D.T., Lambshead, P.J.D., Rogers, A.D., Smerdon, G.R., 2006. Development and evaluation of a DNA barcoding approach for the rapid identification of nematodes. Marine Ecology Progress Series 320, 1–9.
- Bhadury, P., Austen, M.C., Bilton, D.T., Lambshead, P.J.D., Rogers, A.D., Smerdon, G.R., 2008. Evaluation of combined morphological and molecular techniques for marine nematode (*Terschellingia* spp.) identification. Marine Biology 154, 509–518.
- Bhattacharya, B.D., Hwang, J-S., Sarkar, S.K., Rakhsit, D., Murugan, K., Tseng, L-C., 2015. Community structure of mesozooplankton in coastal waters of Sundarban mangrove wetland, India: a multivariate approach. Journal of Marine Systems 141, 112–121.
- Biswas, H., Mukhopadhyay, S.K., De, T.K., Sen, S., Jana, T.K., 2004. Biogenic controls on the air-water carbon dioxide exchange in the Sundarban mangrove environment, northeast coast of Bay of Bengal, India. Limnology Oceanography 49, 95–101.
- Bollasina, M.A., 2014. Hydrology: Probing the monsoon pulse. Nature Climate Change 4, 422–423.
- Bongers, T., 1990. The maturity index: an ecological measure of environmental disturbance based on nematode species composition. Oecologia 83, 14–19.
- Bongers, T., Alkemade, R., Yeates, G.W., 1991. Interpretation of disturbance-induced maturity decrease in marine nematode assemblages by means of the Maturity Index. Marine Ecology Progress Series 76, 135–142.
- Bongers, T., de Goede, R.G.M., Korthals, G.W., Yeates, G.W., 1995. Proposed changes of c-p classification for nematodes. Russian Journal of Nematology 3(1), 61–62.
- Bongers, T., Ferris, H., 1999. Nematode community structure as a bioindicator in environmental monitoring. Trends in Ecology & Evolution 14, 224–228.

- Bongers, T., van der Haar, J., 1990. On the potential of basing an ecological typology of aquatic sediments on the nematode fauna: An example from the river Rhine. Hydrobiological Bulletin 24, 37–45.
- Boonyatumanond, R., Wattayakorn, G., Togo, A., Takada, H., 2006. Distribution and origins of polycyclic aromatic hydrocarbons (PAHs) in riverine, estuarine, and marine sediments in Thailand. Marine Pollution Bulletin 52, 942–956.
- Borja, A., Franco, J., Pe´rez, V., 2000. A marine biotic index to establish the ecological quality of soft bottom benthos within European estuarine and coastal environments. Marine Pollution Bulletin 40, 1100–1114.
- Boucher, G., 1997. Structure and biodiversity of nematode assemblages in the SW lagoon of New Caledonia. Coral Reefs 16, 177–186.
- Boucher, G., Lambshead, P.J.D., 1995. Ecological biodiversity of marine nematodes in samples from temperate, tropical and deep-sea regions. Conservation Biology 9, 1594–1604.
- Boufahja, F., Semprucci, F., Beyrem, H., 2016. An experimental protocol to select nematode species from an entire community using progressive sedimentary enrichment. Ecological Indicators 60, 292–309.
- Bremner, J., Rogers, S.I., Frid, C.L.J., 2003. Assessing functional diversity in marine benthic ecosystems: a comparison of approaches. Marine Ecology Progress Series 254, 11–25.
- Bremner, J., Rogers, S.I., Frid, C.L.J., 2006. Methods for describing ecological functioning of marine benthic assemblages using biological traits analysis (BTA). Ecological Indicators 6, 609–622.
- Broom, M.J., 1982. Structure and seasonality in a Malaysia mudflat community. Estuarine, Coastal and Shelf Science 15, 135–150.
- Buchanan, J.B., 1984. Sediment analysis. In: Holme, N.A., McIntyre, A.D., (Eds.), Methods for the Study of Marine Benthos. Oxford Blackwell Scientific Publications, pp. 41–65.

- Carpentier, A., Como, S., Dupuy, C., Lefrançois, C., Feunteun, E., 2014. Feeding ecology of *Liza* spp. in a tidal flat: evidence of the importance of primary production (biofilm) and associated meiofauna. Journal of Sea Research 92, 86–91.
- Castel, J., 1992. The meiobenthos of coastal lagoon ecosystems and their importance in the food web.Vie et Milieu 42, 125–135.
- Chalcraft, D.R., Resetarits, W.J., 2003. Mapping functional similarity of predators on the basis of trait similarities. American Naturalist 162, 390–402.
- Chanda, A., Das, S., Mukhopadhyay, A., Ghosh, A., Akhand, A., Ghosh, P., Ghosh, T., Mitra, D., Hazra, S., 2018. Sea surface temperature and rainfall anomaly over the Bay of Bengal during the El Niño-Southern Oscillation and the extreme Indian Ocean Dipole events between 2002 and 2016. Remote Sensing Applications: Society and Environment 12, 10–22.
- Chandler, G.T., Fleeger, J.W., 1984. Tube-building by a marine meiobenthic harpacticoid. Chen, C.A., Soo, C.L., Balsamo, M., Semprucci, F., 2018. An approach based on nematode descriptors for the classification of ecological quality (EcoQ) of the Malaysian coasts. Marine Biodiversity 48(1), 117–126.
- Chatterjee, M., Shankar, D., Sen, G.K., Sanyal, P., Sundar, D., Michael, G., Chatterjee, A., Amol, P., Mukherjee, D., Suprit, K., Mukherjee, A., Vijith, V., Chatterjee, S., Basu, A., Das, M., Chakraborti, S., Kalla, A., Misra, S., Mukhopadhyay, S., Mandal, G., Sarkar, K., 2013. Tidal variations in the Sundarbans Estuarine System, India. Journal of Earth System Science 122(4), 899–933.
- Chinnadurai, G., Fernando, O.J., 2003. Meiofauna of Pichavaram mangroves along the southeast coast of India. Journal of the Marine Biological Association of India 45, 158–165.
- Chinnadurai, G., Fernando, O.J., 2006a. New records of free-living marine nematodes from India. Records of Zoological Survey of India 106(Part 4), 45–54.
- Chinnadurai, G., Fernando, O.J., 2006b. Meiobenthos of Cochin mangroves (South west coast of India) with emphasis on free-living marine nematode assemblages. Russian Journal of Nematology 14(2), 127–137.

- Chinnadurai, G., Fernando, O.J., 2007a. Meiofauna of mangroves of the southeast coast of India with special reference to the free-living marine nematode assemblage. Estuarine, Coastal and Shelf Science 72, 329–336.
- Chinnadurai, G., Fernando, O.J., 2007b. Impact of mangrove leaves on meiofauna density: An experimental approach, The ICFAI Journal of Life Sciences 1(1), 62–70.
- Choe, A., von Reuss, S.H., Kogan, D., Gasser, R.B., Platzer, E.G., Schroeder, F.C., Sternberg, P.W., 2012. Ascaroside signaling is widely conserved among nematodes. Current Biology 22(9), 772–780.
- Chowdhury, C., Majumder, N., Ray, R., Jana, T.K., 2012. Inter-annual abundance variation in some genera of diatom and zooplankton in mangrove ecosystem. Biodiversity Conservation 21(8), 2029–2043.
- Clarke, K.R., Gorley, R.N., 2006. PRIMER v6: User manual/tutorial. Plymouth: PRIMER-E.
- Clarke, K.R., Somerfield, P.J., Chapman, M.G., 2008. Testing of null hypotheses in explanatory community analyses: Similarity profiles and biota-environment linkage. Journal of Experimental Marine Biology and Ecology 366, 56–69.
- CMFRI, FRAD. 2017, Marine fish landings in India 2016. Technical Report, ICAR-Central Marine Fisheries Research Institute, Kochi, 10.
- Collins, M., Knutti, R., Arblaster, J.M., Dufresne, J.L., Fichefet, T., Gao, X., Gutowski, W.J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A.J., Wehner, M., 2013. In: Stocker, T., Qin, D., Plattner, G., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P., (Eds.), Long-term Climate Change: Projections, Commitments and Irreversibility. Climate Change 2013: the Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge University Press, Cambridge and New York, pp. 1029–1136.
- Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R.V., Paruelo, J., Raskin, R.G., Sutton, P., van den Belt, M., 1997. The value of the world's ecosystem services and natural capital. Nature 387, 253–260.

- Coull, B.C., 1970. Shallow water meiobenthos of the Bermuda platform. Oecologia (Berl) 4, 325–357.
- Coull, B.C., 1973. Estuarine meiofauna: a review: trophic relationships and microbial interactions. In: Stevenson, H., Colwell, R.R., (Eds), Estuarine Microbial Ecology. University of South Carolina Press, Columbia, pp. 499–512.
- Coull, B.C., 1985. Long-term variability of estuarine meiobenthos: an 11 years study. Marine Ecology Progress Series 24, 205–218.
- Coull, B.C., 1988. Ecology of the marine meiofauna. In: Higgins, R.P., Thiel, H., (Eds.), Introduction to the Study of the Meiofauna. Smithsonian Institution Press, Washington, DC, pp. 18–38.
- Coull, B.C., 1990. Are members of the meiofauna food for higher trophic levels? Transactions of the American Microscopical Society 233–246.
- Coull, B.C., 1999. Role of meiobenthos in estuarine soft-bottom habitats. Australian Journal of Ecology 24, 327–343.
- Coull, B.C., Bell, S.S., 1979. Perspectives of marine meiofaunal ecology. In: Livingston, R.J., (Eds.), Ecological processes in coastal and marine ecosystems, New York, Plenum publishing company, pp. 189–216.
- Coull, B.C., Chandler, G.T., 1992. Pollution and meiofauna: field, laboratory, and mesocosms studies. Oceanography and Marine Biology: An Annual Review 30,191– 271.
- Covich, A.P., Austen, M.C., Bärlocher, F., Chauvet, E., Cardinale, B.J., Biles, C.L., Inchausti, P., Dangles, O., Solan, G.M.O., Statzner, M.B., 2004. The role of biodiversity in the functioning of freshwater and marine benthic ecosystems. Bioscience 54(8), 767–775.
- Cullen, D.J., 1973. Bioturbation of superficial marine sediments by interstitial meiobenthos. Nature 242, 323–324.
- Curini-Galletti, M., Artois, T., Delogu, V., De Smet, W.H., Fontaneto, D., Jondelius, U., Leasi, F., Martinez, A., Meyer-Wachsmuth, I., Nilsson, K.S., Tongiorgi, P., Worsaae,

K., Todaro, M.A., 2012. Patterns of diversity in soft-bodied meiofauna: dispersal ability and body size matter. PLoS One 7, e33801.

- Dal Zotto, M., Santullic, A., Simonini, R., Todaroba, M.A., 2016. Organic enrichment effects on a marine meiofauna community, with focus on Kinorhyncha. Zoologischer Anzeiger 265, 127–140.
- Damodaran, R., 1973. Studies on the benthos of the mud banks of the Kerala Coast. PhD Thesis. Cochin University of Science and Technology, pp. 1–126.
- Danda, A.A., Gayatri, S., Ghosh, A., Bandyopadhyay, J., Hazra, S., 2011. Indian Sundarbans delta: a vision. World Wide Fund for Nature, India.
- Danielsen, F., Sørensen, M.K., Olwig, M.F., Selvam, V., Parish, F., Burgess, N.D., Hiraishi, T., Karunagaran, V.M., Rasmussen, M.S., Hansen, L.B., Quarto, A., Suryadiputra, N., 2005. The Asian tsunami: a protective role for coastal vegetation. Science 310, 643.
- Danovaro, R., Gambi, C., Croce, N.D., 2002. Meiofauna hotspot in the Atacama Trench, eastern South Pacific Ocean. Deep-Sea Research I 843–857.
- Danovaro, R., Dell'Anno, A., Pusceddu, A., 2004a. Biodiversity response to climate change in a warm deep sea. Ecology letters 7, 821–828.
- Danovaro, R., Gambi, C., Mirto, S., Sandulli, R., Ceccherelli, V.U., 2004b. Meiofauna. Biologia Marina Mediterranea 11, 55–97.
- Danovaro, R., Dell'Anno, A., Pusceddu, A., Gambi, C., Heiner, I., Kristensen, R.M., 2010. The first metazoa living in permanently anoxic conditions. BMC Biology 8, 1–30.
- Datta, T.K., Navarrete, A.D.J., Mahapatro, A., 2015. *Rhynchonema dighaensis* sp. nov. (Monhysterida: Xyalidae): a marine nematode from the Indian coast with an illustrated guide and modified key for species of *Rhynchonema* Cobb, 1920. Zootaxa 3905(3), 365–380.
- Dauvin, J.C., 1998. The fine sand *Abra alba* community of the Bay of Morlaix twenty years after the Amoco Cadiz oil spill. Marine Pollution Bulletin 36, 669–676.

- Della Patrona, L., Marchand, C., Hubas, C., Molnar, N., Deborde, J., Meziane, T., 2016. Meiofauna distribution in a mangrove forest exposed to shrimp farm effluents (New Caledonia). Marine Environmental Research 119, 100–113.
- De Troch, M., Gurdebeke, S., Fiers, F., Vincx, M., 2001. Zonation and structuring factors of meiofauna communities in a tropical sea grass bed (Gazi Bay, Kenya). Journal of Sea Research 45, 45–61.
- De Troch, M., Roelofs, M., Riedel, B., Grego, M., 2013. Structural and functional responses of harpacticoid copepods to anoxia in the Northern Adriatic: an experimental approach. Biogeosciences 10, 4259–4272.
- Derycke, S., Remerie, T., Vierstraete, A., Backeljau, T., Vanfleteren, J., Vincx, M., Moens, T., 2005. Mitochondrial DNA variation and cryptic speciation within the free-living marine nematode *Pellioditis marina*. Marine Ecology Progress Series 300, 91–103.
- Derycke, S., Hendrickx, F., Backeljau, T., D'Hondt, S., Camphijn, L., Vincx, M., Moens, T., 2007. Effects of sublethal abiotic stressors on population growth and genetic diversity of *Pellioditis marina* (Nematoda) from the Westerschelde estuary. Aquatic Toxicology 82, 110–119.
- Derycke, S., Vanaverbeke, J., Rigaux, A., Backeljau, T., Moens, T., 2010. Exploring the use of Cytochrome Oxidase c subunit 1 (COI) for DNA barcoding of free-living marine nematodes. PLoS ONE 5(10), e13716.
- Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., Martin, P.R., 2008. Impacts of climate warming on terrestrial ectotherms across latitude. Proceedings of the National Academy of Sciences of the United States of America 105, 6668–6672.
- Dey, M., Ganguly, D., Chowdhury, C., Majumder, N., Jana, T.K., 2012. Intra-Annual Variation of Modern Foraminiferal Assemblage in a Tropical Mangrove Ecosystem. Wetlands 32(5), 813–826.
- Diaz, S., Cabido, M., 2001. Vive la différence: plant functional diversity matters to ecosystem processes. Trends in Ecology & Evolution 16, 646–655.

- Dittmann, S., 1996. Effects of macrobenthic burrows on infaunal communities in tropical tidal flats. Marine Ecology Progress Series 134, 119–130.
- dos Santos, G., Corrêa, G., Valdes, Y., Apolônio Silva de Oliveira, D., Fonsêca-Genevois, V., Silva, A., Pontes, L., Dolan, E., Ingels, J., 2018. *Eretmochelys imbricata* shells present a dynamic substrate for a facilitative epibiont relationship between macrofauna richness and nematode diversity, structure and function. Journal of Experimental Marine Biology and Ecology 502, 153–163.
- Doulgeraki, S., Lampadariou, N., Sinis, A., 2006. Meiofaunal community structure in three Mediterranean coastal lagoons (North Aegean Sea). Journal of the Marine Biological Association of the United Kingdom 86, 209–220.
- Dovgal, I.V., Chatterjee, T., Ingole, B., 2009. New Records of *Thecacineta cothurnioides* and *Trematosoma rotunda* (Ciliophora, Suctorea) as epibionts on nematodes from the Indian Ocean. Protistology 6, 19–23.
- Dujardin, F., 1851. Sur un petit animal marin, l'Echinodère, formant un type intermédiaire entre les Crustacés et les Vers. Ann Sci Nat Zoologie, Sér 3 15, 158–160.
- Dye, A.H., 1983. Vertical and horizontal distribution of meiofauna in mangrove sediments in Transkei, Southern Africa. Estuarine, Coastal and Shelf Science 16(6), 591–598.
- Easterling, D.R., Meehl, G.A., Parmesan, C., Changnon, S.A., Karl, T.R., Mearns, L.O., 2000. Climate extremes: observations, modeling and impacts. Science 289, 2068–2074.
- Edwards, D.P., Tobias, J.A., Sheil, D., Meijaard, E., Laurance, W.F., 2014. Maintaining ecosystem function and services in logged tropical forests. Trends in Ecology and Evolution 29, 511–520.
- van Eenennaam, J.S., Rohal, M., Montagana, P.A., Radović, J.R., Oldenburg, T.B.P., Romerod, I.C., Murke, A.J., Foekemae, E.M., 2019. Ecotoxicological benthic impacts of experimental oil-contaminated marine snow deposition. Marine Pollution Bulletin 141, 164–175.
- Elarbaoui, S., Richard, M., Boufahja, F., Mahmoudi, E., Thomas-Guyon, H., 2015. Effect of crude oil exposure and dispersant application on meiofauna: an intertidal

mesocosm experiment. Environmental Sciences: Processes and Impacts 17(5), 997–1004.

- Elliott, M., Quintino, V., 2007. The estuarine quality paradox, environmental homeostasis and the difficulty of detecting anthropogenic stress in naturally stressed areas. Marine Pollution Bulletin 54, 640–645.
- El Wakeel, S.K., Riley, J.P., 1957. The determination of organic carbon in the marine muds. ICES Journal of Marine Science 22(2), 180–183.
- Fattorini, S., Lorenzo, T.D., Galassi, D.M.P., 2018. Earthquake impacts on microcrustacean communities inhabiting groundwater-fed springs alter speciesabundance distribution patterns. Scientific Reports 8(1), 1501.
- Fegley, S.R., 1987. Experimental variation of near-bottom current speeds and its effects on depth distribution of sand-living meiofauna. Marine Biology 95, 183–191.
- Feller, R.J., Warwick, R.M., 1988. Energetics. In: Higgins, R.P., Thiel, H., (Eds.), Introduction to the Study of Meiofauna. Smithsonian Institution Press, Washington, DC and London, pp. 181–196.
- Fernandez-Leborans, G., Chatterjee, T., Grego, M., 2012. New records of epibiont ciliates (Ciliophora) on Harpacticoida (Copepoda, Crustacea) from the bay of Piran (Gulf of Trieste, Northern Adriatic). Cahiers de Biologie Marine 53, 53–63.
- Ferrero, T.J., Debenham, N.J., Lambshead, P.J.D., 2008. The nematodes of the Thames estuary: assemblage structure and biodiversity, with a test of Attrill'slinear model. Estuarine, Coastal and Shelf Science 79, 409–418.
- Ferris, H., Bongers, T., 2009. Indices developed specifically for nematode assemblages.In: Wilson, M.J., Kakouli-Duarte, T., (Eds.), Nematodes as Environmental Indicators. CABI Publishing, Oxfordshire, UK, pp. 124–145.
- Finch, B.E., Stefansson, E.S., Langdon, C.J., Pargee, S.M., Blunt, S.M., Gage, S.J., Stubblefield, W.A., 2016. Photo-enhanced toxicity of two weathered Macondo crude oils to early life stages of the eastern oyster (*Crassostrea virginica*). Marine Pollution Bulletin 113, 316–323.

- Findlay, S.E.G., 1981. Small scale spatial distribution of meiofauna on a mud and sand flat. Estuarine, Coastal and Shelf Science 12, 471–484.
- Fisher, R., 2003. Ciliate hitch-hikers-nematode ecto-commensals from tropical Australian sea-grass meadows. Journal of the Marine Biological Association of the United Kingdom 83, 445–446.
- Fleeger, J.W., Chandler, G.T., Fitzhugh, G.R. Phillips, F.E., 1984. The effects of tidal currents on meiofauna densities in vegetated salt marsh sediments. Marine Ecology Progress Series 19, 49–53.
- Folk, R.L., 1968. Petrology of sedimentary rocks. Austin, TX: Hemphills.
- Fonseca, G., Derycke, S., Moens, T., 2008. Integrative taxonomy in two free-living nematode species complexes. Biological Journal of the Linnean Society 94, 737– 753.
- Fontaneto, D., Iakovenko, N., De Smet, W.H., 2015. Diversity gradients of rotifer species richness in Antarctica. Hydrobiologia 761(1), 235–248.
- Frontalini, F., Semprucci, F., Armynot du Châtelet, E., Francescangeli, F., Margaritelli, G., Rettori, R., Spagnoli, F., Balsamo, M., Coccioni, R., 2014. Biodiversity trends of the meiofauna and foraminifera assemblages of Lake Varano (southern Italy). Proceedings of the Biological Society of Washington 127, 7–22.
- Frontalini, F., Semprucci, F., Di Bella, L., Caruso, A., Cosentino, C., Maccotta, A., Scopelliti, G., Sbrocca, C., Bucci, C., Balsamo, M., Martins, M.V., du Châtelet, E.A., Coccioni, R., 2018. The response of cultured meiofaunal and benthic foraminiferal communities to lead exposure: Results from mesocosm experiments. Environmental Toxicology and Chemistry 37(9), 2439-2447.
- Gallucci, F., Moens, T., Vanreusel, A., Fonseca, G., 2008. Active colonization of disturbed sediments by deep-sea nematodes: evidence for the patch mosaic model. Marine Ecology Progress Series 367, 173–183.
- Gambi, C., Bianchelli, S., Pérez, M., Invers, O., Ruiz, J., Danovaro, R., 2009. Biodiversity response to experimental induced anoxia in seagrass sediments. Biodiversity and Conservation 18, 33–54.

- Ganapati, P.N., Rao, M.V.L., 1959. Preliminary observations on the bottom fauna of the continental shelf of the northeast coast of India. Proceedings of The First All India Congress of Zoology Part 3, 8–13.
- Gauch, H.G., 1982. Multivarlate Analysis in Community Ecology. Cambridge University Press, New York.
- Gaever, S., Moodley, L., Pasotti, F., Houtekamer, M., Middelburg, J.J., Danovaro, R., Vanreusel, A., 2009. Trophic specialisation of metazoan meiofauna at the Håkon Mosby Mud Volcano: fatty acid biomarker isotope evidence. Marine Biology 156, 1289–1296.
- Gee, J.M., 1987. Impact of epibenthic predation on estuarine intertidal harpacticoid copepod populations. Marine Biology 96, 497–510.
- Gee, J.M., 1989. An ecological and economic review of meiofauna as food for fish. Zoological Journal of the Linnean Society 96(3), 243–261.
- Gerlach, S.A., 1978. Food chain relationship in subtidal silty sand marine sediment and the role of meiofauna in stimulating bacterial productivity. Oecologia 33, 55–69.
- Ghosh, A., Biswas, S., Barman, P., 2014. Marsh foraminiferal assemblages in relation to vegetation in Sunderban, India. Journal of the Geological Society of India 84, 657– 667.
- Ghosh, M., Mandal, S., 2018. Free-living marine nematode diversity from the Indian coasts. Marine Biodiversity 48(1), 179–194.
- Ghosh, M., Mandal, S., Chatterjee, M., 2018. Impact of unusual monsoonal rainfall in structuring meiobenthic assemblages at Sundarban estuarine system, India. Ecological Indicators 94, 139–150.
- Ghosh, M., Mandal, S., 2019a. Living with nematode: an epibiont *Trematosoma rotunda* associated with basibiont *Desmodora scaldensis* from Matla Estuary, Sundarbans, India. Thalassas: An International Journal of Marine Sciences 35, 619–624.
- Ghosh, M., Mandal, S., 2019b. Does vertical distribution of meiobenthic community structure differ among various mangrove habitats of Sundarban Estuarine System? Regional Studies in Marine Science 31, 100778.

- Ghosh, M., Mandal, S., 2021. Deciphering the synergistic impact of elevated temperature and oil pollution on meiobenthic community structure: A benthocosm study. Ecotoxicology and Environmental Safety 207, 111549.
- Giere, O., 2009. Meiobenthology. The microscopic motile fauna of aquatic sediments, 2nd edn. Springer-Verlag, Berlin.
- Gili, J.M., 2002. Towards a transitory or ephemeral key habitat concept. Trends in Ecology & Evolution 17, 453.
- Giménez, L., Borthagaray, A.E., Rodri'guez M., Brazeiro, A., Dimitriadis, C., 2005. Scale-dependent patterns of macrofaunal distribution in soft-sediment intertidal habitats along a large-scale estuarine gradient. Helgoland Marine Research 59, 224– 236.
- Gingold, R., Moens, T., Rocha-Olivares, A., 2013. Assessing the response of nematode communities to climate change-driven warming: a microcosm experiment. PLoS ONE 8(6), e66653.
- Goldin, Q., Mishra, V., Ullal, V., Athalye, R., Gokhale, S., 1996. Meiobenthos of mangrove mudflats from shallow region of Thane creek, central west coast of India. Indian Journal of Marine Sciences 25, 137–141.
- Gopal, B., Chauhan, M., 2006. Biodiversity and its conservation in the Sundarban Mangrove Ecosystem. Aquatic sciences 68, 338–354.
- Guo, Y., Somerfield, P.J., Warwick, R.M., Zhang, Z., 2001. Large-scale patterns in the community structure and biodiversity of free-living nematodes in the Bohai Sea, China. Journal of the Marine Biological Association of the United Kingdom 81, 755–763.
- Gwyther, J., 2000. Meiofauna in phytal-based and sedimentary habitats of a temperate mangrove ecosystem - a preliminary survey. Proceedings of the Royal Society of Victoria 112, 137–151.
- Gwyther, J., 2003. Nematode assemblages from *Avicennia marina* leaf litter in a temperate mangrove forest in south-eastern Australia. Marine Biology 142, 289–297.

- Gyedu-Ababio, T.K., Furstenberg, J.P., Baird, D., Vanreusel, A., 1999. Nematodes as indicators of pollution: a case study from the Swartkops River system, South Africa. Hydrobiologia 397, 155–169.
- Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C.,
  Bruno, J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan,
  H.S., Madin, E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson,
  R., 2008. A global map of human impact on marine ecosystems. Science 319(5865),
  948–952.
- Hamerlynck, O., Vanreusel, A., 1993. *Mesacanthion-Diplechma* (Nematoda, Thoracostomopsidae), a link to higher trophic levels. Journal of the Marine Biological Association of the United Kingdom 73, 453–456.
- Harkantra, S.N., Parulekar, A.H., 1989. Population distribution of meiofauna in relation to some environmental features in a sandy intertidal region of Goa, west coast of India. Indian Journal of Marine Sciences 20, 232–234.
- Harrison, P.J., Khan, N., Yin, K., Saleem, M., Bano, N., Nisa, M., Ahmed, S.I., Rizvi, N., Azam. F., 1997. Nutrient and phytoplankton dynamics in two mangrove tidal creeks of the Indus river delta Pakistan. Marine Ecology Progress Series 157, 13–19.
- Harley, C.D.G., Hughes, A.R., Hultgren, K.M., Miner, B.G., Sorte, C.J.B., Thornber, C.S., Rodriguez, L.F., Tomanek, L., Williams, S.L., 2006. The impacts of climate change in coastal marine systems. Ecology Letters 9, 228–241.
- Heip, C., 1980. Meiobenthos as a tool in the assessment of marine environmental quality.Rapports et Proces-verbaux des Réunions. Conseil International pour l'Éxploration de la Mer 179, 182–187.
- Heip, C., Vincx, M., Vranken, G., 1985. The ecology of marine nematodes. Oceanography and Marine Biology: An Annual Review 23, 399–489.
- Heip, C., Huys, R., Vincx, M., Vanreusel, A., Smol, N., Herman, R. Herman, P.M.J., 1990. Composition, distribution, biomass and production of North Sea meiofauna. Netherlands Journal of Sea Research 26, 333–342.

- Higgins, R.P., Thiel, H., 1988. Introduction to the study of meiofauna. Smithsonian Institution Press.
- Hodda, M., Nicholas, W.L., 1985. Meiofauna Associated with Mangroves in the Hunter River Estuary and Fullerton Cove, South-eastern Australia. Australian Journal of Marine and Freshwater Research 36, 41–50.
- Holbrook, N.J., Sen Gupta, A., Oliver, E.C.J., Hobday, A.J., Benthuysen, J.A., Scannell, H.A., Smale, D.A., Wernberg, T., 2020. Keeping pace with marine heatwaves. Nature Reviews Earth & Environment 1, 482–493.
- Holmer, M., Duarte, C.M., Heilskov, A., Olesen, B., Terrados, J., 2003. Biogeochemical conditions in sediments enriched by organic matter from net-pen fish farms in the Bolinao area, Philippines. Marine Pollution Bulletin 46, 1470–1479.
- Honda, M., Suzuki, N., 2020. Toxicities of polycyclic aromatic hydrocarbons for aquatic animals. International Journal of Environmental Research and Public Health 17, 1363.
- Hubas, C., Sachidhanandam, C., Rybarczyk, H., Lubarsky, H.V., Rigaux, A., Moens, T., Paterson, D.M., 2010. Bacterivorous nematodes stimulate microbial growth and exopolymer production in marine sediment microcosms. Marine Ecology Progress Series 419, 85–94.
- Huys, R., Herman, P.M.J., Heip, C.H.R., Soetaert, K., 1992. Meiobenthos of the North sea, diversity and biomass trends of distribution of copepod community. ICES Journal of Marine Science 49, 23–44.
- Ingels, J., Kiriakoulakis, K., Wolff, G.A., Vanreusel, A., 2009. Nematode diversity and its relation to the quantity and quality of sedimentary organic matter in the deep Nazare' Canyon, Western Iberian Margin. Deep-Sea Research Part I 56, 1521–1539.
- Ingels, J., Dashfield, S.L., Somerfield, P.J., Widdicombe, S., Austen, M.C., 2014. Interactions between multiple large macrofauna species and nematode communities– mechanisms for indirect impacts of trawling disturbance. Journal of Experimental Marine Biology and Ecology 456, 41–49.

- Ingole, B.S., Parulekar, A.H., 1998. Role of salinity in structuring the intertidal meiofauna of a tropical estuarine beach: Field evidence. Indian Journal of Geo-Marine Sciences 27(3–4), 356–361.
- Ingole, B.S., Goltekar, R., 2004. Subtidal micro and meiobenthic community structure in the Gulf of Kachchh. Proceedings of MBR 395–419.
- Ingole, B., Sivadas, S., Goltekar, R., Clemente, S., Nanajkar, M., Sawant, R., D'Silva, C., Sarkar, A., Ansari, Z., 2006. Ecotoxicological effect of grounded MV River Princess on the intertidal benthic organisms off Goa. Environment International 32, 284–291.
- IPCC, 2014. In: Pachauri, R.K., Meyer, L.A., (Eds.), Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Switzerland, Geneva, 151 pp.
- Irizuki, T., Ito, H., Sako, M., Yoshioka, K., Kawano, S., Nomura, R., Tanaka, Y., 2015. Anthropogenic impacts on meiobenthic Ostracoda (Crustacea) in the moderately polluted Kasado Bay, Seto Inland Sea, Japan, over the past 70 years. Marine Pollution Bulletin 91, 149–159.
- Jax, K., 2005. Function and "functioning" in ecology: what does it mean? Oikos 111, 641–648.
- Jensen, P., 1987. Feeding ecology of free-living aquatic nematodes, Marine Ecology Progress Series 35, 87–196.
- Kang, T., Min, W.G., Rho, H.S., Park, H.S., Kim, D., 2014. Differential responses of a benthic meiofaunal community to an artificial oil spill in the intertidal zone. Journal of the Marine Biological Association of the United Kingdom 94(2), 219–231.
- Kang, T., Oh, J., Hong, J., Kim, D., 2016. Effect of the Hebei Spirit Oil Spill on Intertidal Meiofaunal Communities in Taean, Korea. Marine Pollution Bulletin 113, 444–453.
- Kathiresan, K., Bingham, B.L., 2001. Biology of mangroves and mangrove ecosystems. Advances in Marine Biology 40, 81–251.
- Kennedy, A.D., Jacoby, C.A., 1999. Biological indicators of marine environmental health: meiofauna - a neglected benthic component? Environmental Monitoring and Assessment 54(1), 47–68.

- Kennish, M.J., 2002. Environmental threats and environmental future of estuaries. Environmental Conservation 29, 78–107.
- Khan, R.A., 2003. Biodiversity of macrobenthos on the intertidal flats of Sunderban estuarine region, India. Records of the Zoological Survey of India 101(Part 3-4), 181–205.
- Kinne, O., 1977. Environmental factors—salinity: animal—invertebrates. In: Kinne, O., (Eds.), Marine ecology. New York: John Wiley, pp. 821–996.
- Kotal, S.D., Roy, S.S., Bhowmik, S.K.R., 2014. Catastrophic heavy rainfall episode over Uttarakhand during 16–18 June 2013 - observational aspects. Current Science 107(2), 237–245.
- Kovach, W.L., 1998. MVSP-multi-variate statistical package for windows, Ver. 3.0. Kovach computing services, Pentraeth.
- Krishnamurthy, K., Sultan Ali, M.A., Jeyaseelan, M.J.P., 1984. Structure and dynamics of the aquatic food web community with special reference to nematodes in mangrove ecosystem. In: Soepadmo, E., Rao, A.N., MacIntosh, D.J., (Eds.), Proceedings of the Asian Symposium on Mangrove Environment: Research and Management, 1984. University of Malaya, Kuala Lumpur, pp. 429–452.
- Krumhardt, K.M., Lovenduski, N.S., Long, M.C., Lindsay, K., 2017. Avoidable impacts of ocean warming on marine primary production: Insights from the CESM ensembles. Global Biogeochemical Cycles 31,114–33.
- Kumar, A., Sen, D., Bhadury, P., 2015. Unraveling free-living marine nematode community structure from a biodiversity rich tropical coastal setting based on molecular approaches. Marine Biodiversity 45, 537–547.
- Kurien, C.V., 1953. A preliminary survey of the bottom fauna and bottom deposits of the Travancore coast within 15 fathom line. The Proceedings of the National Academy of Sciences, India 19, 746–775.
- Kurien, C.V., 1967. Studies of the benthos of the Southwest coast of India. Bulletin of the National Institute of Sciences of India 38, 649–656.

- Kurien, C.V., 1972. Ecology of benthos in a tropical estuary. The Proceedings of the Indian National Science Academy (B: Biological Sciences) 38(3–4), 156–163.
- Lambshead, P., Platt, H., Shaw, K., 1983. The detection of differences among assemblages of marine benthic species based on an assessment of dominance and diversity. Journal of Natural History 17, 859–874.
- Lambshead, P.J.D., Boucher, G., 2003. Marine nematode deep sea biodiversity– hyperdiverse or hype? Journal of Biogeography 30, 475–485.
- Lambshead, P.J.D., Brown, C.J., Ferrero, T.J., Hawkins, L.E., Smith, C.R., Mitchell, N.J., 2003. Biodiversity of nematodes assemblages from the region off the Clarion– Clipperton Fracture Zone, an area of commercial mining interest. BMC Ecology 3, 1–16.
- Lambshead, P.J.D., 2004. Marine nematode biodiversity. In: Chen, Z.X., Chen, S.Y., Dickson, D.W., (Eds.), Nematology: Advances and perspectives, vol. 1. Wallingford, United Kingdom: CABI Publishing, pp. 439–468.
- Landers, S.C., Nichols, A.C., Barron, N.K., Schimmer, C.A., Tao, R., Yu, K., Stuart, P.M., Ólafsson, E., 2014. Nematode and copepod diversity (2012) from Louisiana near the Deep water Horizon oil spill. Proceedings of the Biological Society of Washington 127(1), 47–57.
- Leduc, D., Rowden, A.A., Pilditch, C.A., Maas, E.W., Probert, P.K., 2013. Is there a link between deep-sea biodiversity and ecosystem function? Marine Ecology 34, 334–344.
- Lee, M.R., Mayorga-Dussarrat, J., 2016. The impact of feeding by Chilean flamingos (*Phoenicopterus chilensis*) on the meiofaunal assemblages of a tidal flat. Marine Biology Research 12(10), 1039–1052.
- Lee, M.R., Torres, R., Manríquez, P.H., 2017. The combined effects of ocean warming and acidification on shallow-water meiofaunal assemblages. Marine Environmental Research 131, 1–9.

- Lejeusne, C., Chevaldonne, P., Pergent-Martini, C., Boudouresque, C.F., Perez, T., 2010. Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea. Trends in Ecology and Evolution 25, 250–260.
- Long, S.M., Ross, O.B.H., 1999. Vertical Distribution of Nematodes (Nematoda) and Harpacticoid Copepods (Copepoda: Harpacticoida) in Muddy and Sandy Bottom of Intertidal Zone of LokKawi, Sabah, Malaysia. The Raffles Bulletin of Zoology 47(2), 349–363.
- Losi, V., Moreno, M., Gaozza, L., Vezzulli, L., Fabiano, M., Albertelli, G., 2013. Nematode biomass and allometric attributes as indicators of environmental quality in a Mediterranean harbour (Ligurian Sea, Italy). Ecological Indicators 30, 80–89.
- Madhu, N.V., Jyothibabu, R., Balachandran, K.K., Honey, U.K., Martin, G.D., Vijay, J.G., Shiyas, C.A., Gupta, G.V.M., Achuthankutty, C.T., 2007. Monsoonal impact on planktonic standing stock and abundance in a tropical estuary (Cochin backwaters— India). Estuarine, Coastal and Shelf Science 73, 54–64.
- Mahmoudi, E., Essid, N., Beyrem, H., Hedfi, A., Boufahja, F., Vitiello, P., Aissa, P., 2005. Effects of hydrocarbon contamination on a free living marine nematode community: results from microcosm experiments. Marine Pollution Bulletin 50, 1197–1204.
- Majdi, N., Threis, I., Traunspurger, W., 2017. It's the little things that count: meiofaunal density and production in the sediment of two headwater streams. Limnology Oceanography 62, 151–163.
- Mandal, S., Ray, S., Ghosh, P.B., 2012. Modelling the impact of mangroves on fish population dynamics of Hooghly—Matla estuarine system, West Bengal, India. Procedia Environmental Sciences 13, 414–444.
- Manini, E., Gambi, C., Danovaro, R., Fabiano, M., 2000. Meiobenthic grazing rates on bacteria and microphytobenthos in the northern Adriatic Sea: Preliminary results. Biolagia Marina Mediterranea 7(1), 233–238.
- Manna, S., Chaudhuri, K., Bhattacharyya, S., Bhattacharyya, M., 2010. Dynamics of Sundarban estuarine ecosystem: Eutrophication induced threat to mangroves. Saline Systems 6, 8.

- Mare, M.F., 1942. A study on marine benthic community with special reference to the micro-organisms. Journal of the Marine Biological Association of the United Kingdom 25, 517–554.
- Maurer, D., 2000. The dark side of the taxonomic sufficiency TS. Marine Pollution Bulletin 40, 98–101.
- McIntyre, A.D., 1968. The meiofauna and macrofauna of some tropical beaches. Journal of Zoology, London 155, 377–392.
- McIntyre, A.D., 1969. Ecology of marine meiobenthos. Biological Reviews 44, 245–290.
- McIntyre, A.D., 1977. Effect of pollution on inshore benthos. In: Coull, B.C., (Eds.), Ecology of marine benthos. (University of South Carolina Press), pp. 301–312.
- McIntyre-Wressnig, A., Bernhard, J.M., McCorkle, D.C., Hallock, P., 2013. Non-lethal effects of ocean acidification on the symbiont-bearing benthic foraminiferal species, Biogeosciences Discuss 8, 9165–9200.
- McLachlan, A., 1978. A quantitative analysis of the meiofauna and the chemistry of the redox potential discontinuity zone in a sheltered sandy beach. Estuarine and Coastal Marine Science 7, 275–290.
- Meadows, A.S., Ingels, J., Widdicombe, S., Hale, R., Rundle, S.D., 2015. Effects of elevated CO<sub>2</sub> and temperature on an intertidal meiobenthic community. Journal of Experimental Marine Biology and Ecology 469, 44–56.
- Mevenkamp, L., Ong, E.Z., Van Colen, C., Vanreusel, A., Guilini, K., 2018. Combined, short-term exposure to reduced seawater pH and elevated temperature induces community shifts in an intertidal meiobenthic assemblage. Marine Environmental Research 133, 32–44.
- Michiels, I., Traunspurger, W., 2004. A three years study of seasonal dynamics of a zoobenthos community in a eutrophic lake. Nematology 6(5), 655–669.
- Miller, D.C., Geider, R.J., Macintyre, H.L., 1996. Microphytobenthos: The ecological role of the secret garden of unvegetated, shallow-water marine habitats. II. Role in sediment stability and shallow-water food webs. Estuaries 19(2A), 202–212.

- Modig, H., Ólafsson, E., 1998. Responses of marine benthic invertebrates to hypoxic events in the Baltic Sea. Journal of Experimental Marine Biology and Ecology 229, 133–148.
- Moens, T., Verbeeck, L., Vincx, M., 1999. Feeding biology of a predatory and a facultatively predatory marine nematode (*Enoploides longispiculosus* and *Adoncholaimus fuscus*). Marine Biology 134, 585–593.
- Moens, T., Vincx, M., 1997. Observations on the feeding ecology of estuarine nematodes. Journal of the Marine Biological Association of the United Kingdom 77, 211–227.
- Moens, T., Vincx, M., 2000. Temperature and salinity constraints on the life cycle of two brackish-water nematode species. Journal of Experimental Marine Biology and Ecology 243, 115–135.
- Mokievsky, V.O., Tchesunov, A.V., Udalov, A.A., Toan, N.D., 2011. Quantitative distribution of meiobenthos and the structure of the free-living nematode community of the mangrove intertidal zone in Nha Trang Bay (Vietnam) in the South China Sea. Russian Journal of Marine Biology 37(4), 272–283.
- Montagna, P.A., Blanchard, F.G., Dinet, A., 1995. Effect of production and biomass of intertidal microphytobenthos on meiofaunal grazing rates. Journal of Experimental Marine Biology and Ecology 185, 149–165.
- Montagna, P.A., Baguley, J.G., Cooksey, C., Hartwell, I., Hyde, L.J., Hyland, J.L., Kalke, R.D., Kracker, L.M., Reuscher, M., Rhodes, A.C., 2013. Deep-sea benthic footprint of the Deepwater Horizon blowout. PLoS One 8(8), e70540.
- Moodley, L., Chen, G.T., Heip, C., Vincx, M., 2000. Vertical distribution of meiofauna in sediments from contrasting sites in the Adriatic Sea: clues to the role of abiotic versus biotic control. Ophelia 53, 203–212.
- Moreno, M., Ferrero, T.J., Gallizia, I., Vezzulli, L., Albertelli, G., Fabiano, M., 2008a. An assessment of the spatial heterogeneity of environmental disturbance within an enclosed harbor through the analysis of meiofauna and nematode assemblages. Estuarine, Coastal and Shelf Science 77, 565–576.

- Moreno, M., Vezzulli, L., Marin, V., Laconi, P., Albertelli, C., Fabiano, M., 2008b. The use of meiofauna diversity as an indicator of pollution in harbours. ICES Journal of Marine Science 65, 1428–1435.
- Moreno, M., Semprucci, F., Vezzulli, L., Balsamo, M., Fabiano, M., Albertelli, G., 2011. The use of nematodes in assessing ecological quality status in the Mediterranean coastal ecosystems. Ecological Indicators 11, 328–336.
- Mortensen, T., 1925. An apparatus for catching the micro-fauna of the sea bottom. Vidensk Medd Dansk Naturh Foren København 80, 445–451.
- Mouillot, D., Graham, N.A.J., Ville'ger. S., Mason, N.W.H., Bellwood, D.R., 2013. A functional approach reveals community responses to disturbances. Trends in Ecology & Evolution 28, 167–177.
- Nanajkar, M.R., Ingole, B.S., 2007. Nematode species diversity as indicator of stressed benthic environment along the central west coast of India. In: Desai, P.V., Roy, R., (Eds.), Diversity and life processes from ocean and land. Goa University, India, pp. 42–52.
- Nanajkar, M.R., Ingole, B.S., 2010. Comparison of tropical nematode communities from the three harbours, west coast of India. Cahiers de Biologie Marine 51(1), 9–18.
- Nandi, S., Choudhury, A., 1983. Quanitative studies on the benthic macrofauna of Sagar Island, intertidal zones, Sundarbans, India. Mahasagar-Bulletin of the National Institute of Oceanography 16(3), 409–414.
- Nandy, T., Mandal, S., Deb, S., Ghosh, M., Nath, T., Chatterjee, M., 2018. Short-term variations in surface water properties in the Sundarban Estuarine System, India. Sustainable Water Resources Management 4(3), 559–566.
- Nandy, T., Mandal, S., Chatterjee, M., 2018. Intra-monsoonal variation of zooplankton population in the Sundarbans Estuarine System, India. Environmental Monitoring and Assessment 190, 603.
- Nascimento, F.J., Karlson, A.M., Näslund, J., Elmgren, R., 2011. Diversity of larger consumers enhances interference competition effects on smaller competitors. Oecologia 166(2), 337–347.

- Nascimento, F.J.A., Naslund, J., Elmgren, R., 2012. Meiofauna enhances organic matter mineralization in soft sediment ecosystems. Limnology and Oceanography 57, 338– 346.
- Nehring, S., 1993. Tube-dwelling meiofauna in marine sediments. International Review of Hydrobiology 78, 521–534.
- Neira, C., Sellanes, J., Soto, A., Gutiérrez, D., Gallardo, V.A., 2001. Meiofauna and sedimentary organic matter off Central Chile: response to changes caused by the 1997–1998 El Niño. Oceanologica Acta 24, 313–328.
- Neres, P.F., Da Fonseca-Genevois, V.G., Torres, R.A., Da Fonseca Cavalcanti, M., De Castro, F.J.V., Da Silva, N.R.R., Rieger, T.T., Decraemer, W., 2010. Morphological and molecular taxonomy of a new *Daptonema* (Nematoda, Xyalidae) with comments on the systematics of some related taxa. Zoological Journal of the Linnean Society 158, 1–15.
- Netto, S.A., Warwick, R.M., Attrill, M.J., 1999. Meiobenthic and macrobenthic community structure in carbonate sediments of Rocas Atoll (North-east Brazil). Estuarine, Coastal and Shelf Science 48, 39–50.
- Netto, S.A., Gallucci, F., 2003. Meiofauna and macrofauna communities in a mangrove from the Island of Santa Catarina, South Brazil. Hydrobiologia 505, 159–170.
- Newell, R.C., 1982. The energetics of detritus utilization in coastal lagoons and near shore waters. In: Oceanologica Acta, Proceedings of the International Symposium on Coastal Lagoons, SCOR/IABO/UNESCO, 8–14 September, 1981, Bordeaux, France, pp. 347–355.
- Nicholas, W.L., Elek, J.A., Stewart, A.C., Marples, T.G., 1991. The nematode fauna of a temperate Australian mangrove mudflat: its population density, diversity and distribution. Hydrobiologia 209, 13–27.
- Nigam, R., Kurtarkar, S.R., Saraswat, R., Linshy, V.N., Rana, S.S., 2008. Response of benthic foraminifera *Rosalina leei* to different temperature and salinity, under laboratory culture experiment. Journal of the Marine Biological Association of the United Kingdom 88, 699–704.

- Nigam, R., Linshy, V.N., Kurtarkar, S.R., Saraswat, R., 2009. Effects of sudden stress due to heavy metal mercury on benthic foraminifer *Rosalina leei*: laboratory culture experiment. Marine Pollution Bulletin 59, 362–368.
- Nozais, C., Perissinotto, R., Tita, G., 2005. Seasonal dynamics of meiofauna in a South African temporarily open/closed estuary (Mdloti Estuary, Indian Ocean). Estuarine, Coastal and Shelf Science 62, 325–338.
- Olafsson, E., 1995. Meiobenthos in mangrove areas in eastern Africa with assemblage structure of free-living marine nematodes, Hydrobiologia 312, 47–57.
- Olafsson, E., Carlstrom, S., Ndaro, S.G.M., 2000. Meiobenthos of hyper saline tropical mangrove sediment in relation to spring tide inundation. Hydrobiologia 426, 57–64.
- Pachauri, R.K., Meyer, L., Van Ypersele, J.P., Brinkman, S., Van Kesteren, L., Leprince-Ringuet, N., Van Boxmeer, F., 2014. Climate Change 2014 Synthesis Report (Geneva, Switzerland).
- Padmakumar, K.B., Thomas, L.C., Vijayan, A., Sanjeevan, V.N., 2014. Occurrence of epibiontic suctorian protozoansa on marine ostracod *Cypridina dentate* (Muller, 1906) in the north eastern Arabian Sea. Indian Journal of Geo-Marine Sciences 44, 1–5.
- Palmer, M.A., 1988. Dispersal of marine meiofauna, a review and conceptual model explaining passive transport and active emergence with implications for recruitment. Marine Ecology Progress Series 48, 81–91.
- Panikkar, N.K., Aiyar, R.C. 1937. The brackish water fauna of Madras. Proceedings of Indian Academy Science 6, 284–337.
- Papa, F., Durand, F., Rossow, W.B., Rahman, A., Bala, S.K., 2010. Satellite altimeterderived monthly discharge of the Ganga-Brahmaputra River and its seasonal to interannual variations from 1993 to 2008. Journal of Geophysical Research 115, C12013.
- Parsons, T.R., Takahashi, M., Hargrave, B., 1990. Biological oceanographic processes. 3rd edition. Oxford: Pergamon Press.

- Parulekar, A.H., Nair, S.A., Harkantra, S.N., Ansari, Z.A., 1976. Some quantitative studies on the benthos of Bombay. Mahasagar 9(1&2), 51–56.
- Pearson, T.H., Black, K.D., 2001. The environmental impacts of marine fish cage culture. In: Black, K.D., (Eds.), Environmental Impacts of Aquaculture. Sheffield Academic Press, Sheffield, pp. 1–31.
- Perugini, M., Visciano, P., Giammarino, A., Manera, M., Di Nardo, W., Amorena, M., 2007. Polycyclic aromatic hydrocarbons in marine organisms from the Adriatic Sea, Italy. Chemosphere 66(10), 1904–1910.
- Petersen, C.G.J., 1913. Havets bonitering. II. Om havbundens dyresamfund og om disses betydning for den marine zoogeografi. Beretn Landbrugsminist Dansk Biolog Station 21, 1–42 plus add 41–68.
- Pfannkuche, O., Thiel, H., 1988. Sample processing. In: Higgins, R.P., Thiel, H., (Eds.), Introduction to the study of Meiofauna, Smithsonian Institute Press, Washington, DC, pp. 134–145.
- Phillips, F.E., Fleeger, J.W., 1985. Meiofauna meso-scale variability in two estuarine habitats. Estuarine, Coastal and Shelf Science 21, 745–756.
- Pielou, E.C., 1969. An Introduction to Mathematical Ecology. Wiley-Interscience, New York, 286 pp.
- Pihl, L., Rosenberg, R., 1984. Food selection and consumption of the shrimp *Crangon crangon* in some shallow marine areas in western Sweden. Marine Ecology Progress Series 15, 159–168.
- Pinto, T.K., Austen, M.C., Warwick, R.M., Somerfield, P.J., Esteves, A.M., Castro, F.J.V., Fonseca-Genevois, V.G., Santos, P.J.P., 2013. Nematode diversity in different microhabitats in a mangrove region. Marine Ecology 34(3), 257–268.
- Piot, A., Nozais, C., Archambault, P., 2014. Meiofauna affect the macrobenthic biodiversity ecosystem functioning relationship. Oikos 123(2), 203–213.
- Platt, H.M., Warwick, R.M., 1980. The significance of free-living nematodes to the littoral ecosystem. In: Price, J.H., Irvine, D.E.G., Farnham, W.F., (Eds), The shore environment, 2nd edn. Academic Press, London, pp. 729–759.

- Platt, H.M., Warwick, R.M., 1983. Free living marine nematodes, Part I, British Enoplids, Pictorial key to world genera and notes for the identification of British species. In: Kermack, D.M., Barnes, R.S.K., (Eds.), Synopses of the British Fauna, vol 28. Cambridge University Press, Cambridge, pp. 1–307.
- Platt, H.M., Warwick, R.M., 1988. Free living marine nematodes, Part II, British Chromodorids, Pictorial key to world genera and notes for the identification of British species. In: Kermack, D.M., Barnes, R.S.K., (Eds.), Synopses of the British Fauna (new series), vol 38. Brill/Backhuys, Leiden, pp. 1–502.
- Poloczanska, E.S., Burrows, M.T., Brown, C.J., GarcõÂa Molinos, J., Halpern, B.S., Hoegh-Guldberg, O., Kappel, C.V., Moore, P.J., Richardson, A.J., Schoeman, D.S., Sydeman, W.J., 2016. Responses of Marine Organisms to Climate Change across Oceans. Frontiers in Marine Science 3, 62.
- Prandle, D., 2009. Estuaries: Dynamics, Mixing, Sedimentation and Morphology. Cambridge University Press, UK.
- Przeslawski, R., Ahyong, S., Byrne, M., Wo"rheide, G., Hutchings, P., 2008. Beyond corals and fish: the effects of climate change on noncoral benthic invertebrates of tropical reefs. Global Change Biology 14, 2773–2795.
- Pusceddu, A., Gambi, C., Manini, E., Danovaro, R., 2007. Trophic state, ecosystem efficiency and biodiversity of transitional aquatic ecosystems: analysis of environmental quality based on different benthic indicators. Chemistry and Ecology 23, 505–515.
- Pusceddu, A., Bianchelli, S., Gambi, C., Danovaro, R., 2011. Assessment of benthic trophic status of marine coastal ecosystems: significance of meiofaunal rare taxa. Estuarine, Coastal and Shelf Science 93, 420–430.
- Pusceddu, A., Bianchelli, S., Martín, J., Puig, P., Palanques, A., Masquéd, P., Danovaro, R., 2014. Chronic and intensive bottom trawling impairs deep-sea biodiversity and ecosystem functioning. Proceedings of the National Academy of Sciences of the United States of America 111, 8861–8866.

- Radwell, A.J., Brown, A.V., 2008. Benthic meiofauna assemblage structure of headwater streams: density and distribution of taxa relative to substrate size. Aquatic Ecology 42(3), 405–414.
- Raes, M., De Troch, M., Ndaro, S.G.M., Muthumbi, A., Guilini, K., Vanreusel, A., 2007. The structuring role of microhabitat type in coral degradation zones: a case study with marine nematodes from Kenya and Zanzibar. Coral Reefs 26, 113–126.
- Rao, G.C., 1980. On the zoogeography of the interstitial meiofauna of the Andaman and Nicobar Islands, Indian Ocean. Records of the Zoological Survey of India 77, 153– 178.
- Rao, G.C., Misra, A., 1983. Studies on the meiofauna of Sagar Island. Proceedings of the Indian Academy of Sciences (Animal Sciences) 92(1), 73–85.
- Rao, G.C., 1986. Meiofauna of the mangrove sediment in South Andaman. Journal of the Andaman Science Association 2(2), 23–32.
- Rao, D.G., 1987a. Ecology of meiobenthos of Rambha Bay in Chilka lagoon, Bay of Bengal. Journal of Marine Biological Association of India 29(1-2), 74–85.
- Rao, G.C., 1987b. Meiofauna of the marine National Park, South Andaman. Journal of the Andaman Science Association 3, 88–97.
- Rao, G.C., 1988. Meiofauna of the intertidal sediments on Great Nicobar. Journal of the Andaman Science Association 4(2), 89–100.
- Rao, G.C., 1989. Intertidal Meiofauna. State Fauna Ser. 1: Fauna of Orissa: Zoological Survey of India, Part-2, pp. 277–318.
- Rao, G.C., 1991. Meiofauna. State Fauna Ser. 2: Fauna of Lakshadweep: Zoological Survey of India, pp. 41–135.
- Rao, G.C., Misra, A., 1983. Studies on the meiofauna of Sagar Island. Proceedings of the Indian Academy of Sciences (Animal Sciences) 92(1), 73–85.
- Reil, L.A., Faubel, A., 1980. Uptake of organic matter by meiofauna organisms and interrelationships with bacteria. Marine Ecology Progress Series 3, 251–256.

- Richardson, A.J., Schoeman, D.S., 2004. Climate impact on plankton ecosystems in the Northeast Atlantic. Science 305(5690), 1609–1612.
- Riemann, F., 1974. On hemisessile nematodes with flagilliform tails living in marine soft bottoms and on microtubes found in deep sea sediments. Mikrofauna Meeresbod 40, 1–15.
- Ristau, K., Spann, N., Traunspurger, W., 2015. Species and trait compositions of fresh water nematodes as indicative descriptors of lake eutrophication. Ecological Indicators 53, 196–205.
- Ritter, A.N., 2012. Effect of eutrophication on benthic microalgae. Master thesis, Department of Biology, Middlebury College, Vermont, 51p.
- Rosenzweig, C., Karoly, D., Vicarelli, M., Neofotis, P., Wu, Q.G., Casassa, G., Menzel, A., Root, T.L., Estrella, N., Seguin, B., Tryjanowski, P., Liu, C.Z., Rawlins, S., Imeson, A., 2008. Attributing physical and biological impacts to anthropogenic climate change. Nature 453, 353–357.
- Ruiz, F., Abad, M., Bodergat, A.M., Carbonel, P., Rodriguez-Lazaro, J., Yasuhara, M., 2005. Marine and brackish-water ostracods as sentinels of anthropogenic impacts. Earth-Science Reviews 72, 89–111.
- Rundle, S.D., Bilton, D.T., Shiozawa, D.K., 2000. Global and regional patterns in lotic meiofauna. Freshwater Biology 44, 123–134.
- Rzeznik-Orignac, J., Fichet, D., Boucher, G., 2003. Spatio-temporal structure of the nematode assemblages of the Brouage mudflat (Marennes Ole´ron, France). Estuarine, Coastal and Shelf Science 58, 77–88.
- Sahoo, G., Suchiang, S.R., Ansari, Z.A., 2013. Meiofauna-mangrove interaction: A pilot study from a tropical mangrove habitat. Cahiers de Biologie Marine 54, 349–358.
- Saikranthi, K., Radhakrishna, B., Rao, T.N., Satheesh, S.K., 2019. Differences in the association of sea surface temperature–precipitating systems over the Bay of Bengal and the Arabian Sea during southwest monsoon season. International Journal of Climatology 39(11), 4305–4312.

- Sajan, S., Damodaran, R., 2007. Faunal composition of meiobenthos from the shelf regions off the west coast of India. Journal of the Marine Biological Association of India 49(1), 19–26.
- Sajan, S., Joydas, T.V., Damodaran, R., 2010. Meiofauna of the western continental shelf of India, Arabian Sea. Estuarine, Coastal and Shelf Science 86, 665–674.
- Samanta, D., Hameed, S. N., Jin, D., Thilakan, V., Ganai, M., Rao, S. A., Deshpande, M., 2018. Impact of a narrow coastal Bay of Bengal sea surface temperature front on an Indian summer monsoon simulation. Scientific Reports 8(1), 17694.
- Samuel, M., 1944. Preliminary observations on the animal communities of the level sea bottom of the Madras coast. Journal of Madras University 15(2), 45–71.
- Sanders, H.L., 1968. Marine benthic diversity: A comparative study. The American Naturalist 102(925), 243–282.
- Sanduli, R., De Nicola, M., 1991. Responses of meiobenthic communities along a gradient of sewage pollution. Marine Pollution Bulletin 22, 463–467.
- Sandulli, R., Semprucci, F., Balsamo, M., 2014. Taxonomic and functional biodiversity variations of meiobenthic and nematode assemblages across an extreme environment: a study case in a blue hole cave. Italian Journal of Zoology 81, 508–516.
- Santos, G.A.P., Derycke, S., Fonse<sup>c</sup>ca-Genevois, V., Coelho, L., Correia, M., Moens, T., 2008. Differential effects of food availability on population growth and fitness of three species of estuarine, bacterial-feeding nematodes. Journal of Experimental Marine Biology and Ecology 355, 27–40.
- Saraswat, R., Kurtarkar, S.R., Mazumder, A., Nigam, R., 2004. Foraminifers as indicators of marine pollution: a culture experiment with *Rosalina leei*. Marine Pollution Bulletin 48, 91–96.
- Saraswat, R., Kouthanker, M., Kurtarkar, S.R., Nigam, R., Naqvi, S.W.A., Linshy, V.N., 2015. Effect of salinity induced pH/alkalinity changes on benthic foraminifera: A laboratory culture experiment. Estuarine, Coastal and Shelf Science 153, 96–107.

- Sarma, A.L.N., Rao, D.G., 1980. The meiofauna of Chilka Lake (Brackish water lagoon). Current Science 49(22), 870–872.
- Sarma, A.L.N., Satpathy, S., Rao, D.G., 1981. Phytal macro and meiofauna of Chilka Lake. Indian Journal of Marine Sciences 10(1), 61–65.
- Sarma, A.L.N., Wilsanand, V., 1994. Littoral meiofauna of Bhitarkanika mangroves of river Mahanadi system, East coast of India. Indian Journal of Marine Sciences 23, 221–224.
- Schneider, K.R., 2008. Heat stress in the intertidal: comparing survival and growth of an invasive and native mussel under a variety of thermal conditions. Biological Bulletin 215, 253–264.
- Schratzberger, M., Warwick, R.M., 1999a. Impact of predation and sediment disturbance by *Carcinus maenas* (L.) on free-living nematode community structure. Journal of Experimental Marine Biology and Ecology 235(2), 255–271.
- Schratzberger, M., Warwick, R.M., 1999b. Differential effects of various types of disturbances on the structure of nematode assemblages: an experimental approach. Marine Ecology Progress Series 181, 227–236.
- Schratzberger, M., Gee, J.M., Rees, H.L., Boyd, S.E., Wall, C.M., 2000. The structure and taxonomic composition of sublittoral meiofauna assemblages as an indicator of the status of the marine environment. Journal of the Marine Biological Association of the United Kingdom 80, 969–980.
- Schratzberger, M., Dinmore, T.A., Jennings, S., 2002. Impacts of trawling on the diversity, biomass and structure of meiofauna assemblages. Marine Biology 140, 83–93.
- Schratzberger, M., Danie, F., Wall, C.M., Kilbride, R., Macnaughton, S.J., Boyd, S.E., Rees, H.L., Swannell, R.P.J., 2003. Response of estuarine meio and macrofauna to *in situ* bioremediation of oil contaminated sediment. Marine Pollution Bulletin 46, 430–443.

- Schratzberger, M., Whomersley, P., Warr, K., Bolam, S.G., Rees, H.L., 2004. Colonisation of various types of sediment by estuarine nematodes via lateral infaunal migration: a laboratory study. Marine Biology 145, 69–78.
- Schratzberger, M., Bolam, S., Whomersley, P., Warr, K., 2006a. Differential response of nematode colonist communities to the intertidal placement of dredged material. Journal of Experimental Marine Biology and Ecology 334, 244–255.
- Schratzberger, M., Warr, K., Rogers, S.I., 2006b. Patterns of nematode populations in the southwestern North Sea and their link to other components of the benthic fauna. Journal of Sea Research 55, 113–127.
- Schratzberger, M., Warr, K., Rogers, S.I., 2007. Functional diversity of nematode community in the southwestern North Sea. Marine Environmental Research 63, 368–389.
- Schratzberger, M., Ingels, J., 2018. Meiofauna matters: the roles of meiofauna in benthic ecosystems. Journal of Experimental Marine Biology and Ecology 502, 12–25.
- Schratzberger, M., Somerfield, P.J., 2020. Effects of widespread human disturbances in the marine environment suggest a new agenda for meiofauna research is needed. Science of the Total Environment 728, 138435.
- Schückel, S., Sell, A.F., Kihara, T.C., Koeppen, A., Kröncke, I., Reiss, H., 2013. Meiofauna as food source for small-sized demersal fish in the southern North Sea. Helgoland Marine Research 67(2), 203–218.
- SCOR Working Group 76, 1994. Suggested criteria for describing deep-sea benthic communities; the final report of SCOR Working Group 76. Progress in Oceanography 34, 81–100.
- Seinhorst, J.W., 1959. A rapid method for the transfer of nematodes from fixative to anhydrous glycerine. Nematologica 4, 67–69.
- Semprucci, F., Colantoni, P., Baldelli, G., Rocchi, M., Balsamo, M., 2010. The distribution of meiofauna on back-reef sandy platforms in the Maldives (Indian Ocean). Marine Ecology 31, 592–607.

- Semprucci, F., Colantoni, P., Sbrocca, C., Baldelli, G., Rocchi, M., Balsamo, M., 2011. Meiofauna in sandy back-reef platforms differently exposed to the monsoons in the Maldives (Indian Ocean). Journal of Marine Systems 87, 208–215.
- Semprucci, F., Balsamo, M., 2012. Key role of free-living nematodes in the marine ecosystem. In: Boeri, F., Jordan, A.C., (Eds.), Nematodes: Morphology, Functions and Management Strategies. NOVA Science Publ., Inc., Hauppauge, pp. 109–134.
- Semprucci, F., Balsamo, M., Frontalini, F., 2014a. The nematode assemblage of a coastal lagoon (Lake Varano, Southern Italy): ecology and biodiversity patterns. Scientia Marina 78, 579–588.
- Semprucci, F., Balsamo, M., 2014b. Free-living Marine Nematodes as Bioindicators: Past, Present and Future Perspectives. Research Trends in Environmental Science 6, 17– 36.
- Semprucci, F., Sbrocca, C., Rocchi, M., Balsamo, M., 2015a. Temporal changes of the meiofaunal assemblage as a tool for the assessment of the ecological quality status. Journal of the Marine Biological Association of the United Kingdom 95, 247–254.
- Semprucci, F., Losi, V., Moreno, M., 2015b. A review of Italian research on free-living marine nematodes and the future perspectives on their use as ecological indicators (Eco Inds). Mediterranean Marine Science 16, 352–365.
- Semprucci, F., Balsamo, M., Sandulli R., 2016. Assessment of the Ecological quality (EcoQ) of the Venice lagoon using the structure and biodiversity of the meiofaunal assemblages. Ecological Indicators 67C, 451-457.
- Semprucci, F., Cesaroni, L., Guidi, L., Balsamo, M., 2018. Do the morphological and functional traits of free-living marine nematodes mirror taxonomical diversity? Marine Environmental Research 135, 114–122.
- Semprucci, F., Facca, C., Ferrigno, F., Balsamo, M., Sfriso, A., Sandulli, R., 2019. Biotic and abiotic factors affecting seasonal and spatial distribution of meiofauna and macrophytobenthos in transitional coastal waters. Estuarine, Coastal and Shelf Science 219, 328–340.

- Sen, A., Ghosh, M., Khanderao, P., Das, S.K., Roy Chowdhury, D., Sarkar, P.K., Saha, R., Bhadury, P., 2016. Modern benthic foraminiferal assemblages from the world's largest deltaic mangrove ecosystem, the Sundarbans. Marine Biodiversity 46, 421– 431.
- Seshappa, G., 1953. Observations on the physical and biological features of the inshore sea bottom along the Malabar Coast. The Proceedings of the National Institute of Sciences of India 19, 257–279.
- Shannon, C., Weaver, W., 1949. The Mathematical Theory of Communication. University of Illinois Press, Urbana, 121 pp.
- Shirayama, Y., Ohta, S., 1990. Meiofauna in a cold deep community of Hatsushima. Central Japan. Journal of the Oceanographic Society of Japan 46, 118–124.
- Simpson, E.H., 1949. Measurement of diversity. Nature 163, 688.
- Singh, R., Ingole, B.S., 2016. Structure and function of nematode communities across the Indian western continental margin and its oxygen minimum zone. Biogeosciences 13, 191–209.
- Singh, R., Ingole, B.S., Nanajkar, M.R., 2009. The life cycle of the free-living marine nematode *Innocuonema tentabunda* de man, 1890. Nematologia Mediterranea 37, 235–238.
- Sinha, B., Choudhury, A., Baqri, G.H., 1987. Studies on the nematodes from mangrove swamps of deltaic Sundarbans, West Bengal, India. III. *Anoplostoma macrospiculum* N. sp. (Anoplostomatidae: Nematoda). Current Science 56(11), 539–540.
- Snelgrove, P.V.R., Blackburn, T.H., Hutchings, P.A.A., Alongi, D.M., Grassle, J.F., Hummel, H., King, G., Koike, I., Lambshead, P.J.D., Ramsing, N.D., Solis-Weiss, V., 1997. The importance of marine sediment biodiversity in ecosystem processes. Ambio: A Journal of the Human Environment 26, 578–584.
- Snelgrove, P.V.R., 1999. Getting to the bottom of marine biodiversity: Sedimentary habitats. BioScience 49, 129–138.

- Soetaert, K., Vincx, M., Wittoeck, J., Tulkens, M., 1995. Meiobenthic distribution and nematode community structure in five European estuaries. Hydrobiologia 311, 185–206.
- Soetaert, K., Muthumbi, A., Heip, C., 2002. Size and shape of ocean margin nematodes: morphological diversity and depth-related patterns. Marine Ecology Progress Series 242, 179–193.
- Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L., 2007. The physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. (Eds.), Cambridge: Cambridge University Press.
- Somerfield, P.J., Clarke, K.R., 1995. Taxonomic levels, in marine community studies revisited. Marine Ecology Progress Series 127, 113–119.
- Somerfield, P.J., Gee, J.M., Aryuthaka, C., 1998. Meiofaunal communities in a Malaysian mangrove forest. Journal of the Marine Biological Association of the United Kingdom 78, 717–732.
- Somerfield, P.J., Yodnarasri, S., Aryuthaka, C., 2000. The effects of green mussel *Perna* viridis (L.) (Mollusca: Mytilidae) culture on nematode community structure in the Gulf of Thailand. The Raffles Bulletin of Zoology 48, 263–272.
- Spalding, M.D., Blasco, E., Field, C.D., 1997. World Mangrove Atlas. International Society for Mangrove Ecosystems, Okinawa, Japan.
- Spalding, M.D., Kainuma, M., Collins, L., 2010. World Atlas of Mangroves. Earthscan, London, 304 pp.
- Steyaert, M., Garner, N., Van Gansbeke, D., Vincx, M., 1999. Nematode communities from the North Sea: environmental controls on species diversity and vertical distribution within the sediment. Journal of the Marine Biological Association of the United Kingdom 79, 253–264.
- Steyaert, M., Vanaberbeke, J., Vanreusel, A., Barranguet, C., Lucas, C., Vincx, M., 2003. The importance of fine-scale, vertical profiles in characterizing nematode community structure. Estuarine, Coastal and Shelf Science 58, 353–366.

- Steyaert, M., Moodley, L., Nadong, T., Moens, T., Soetaert, K., Vincx, M., 2007. Responses of intertidal nematodes to short-term anoxic events. Journal of Experimental Marine Biology and Ecology 345, 175–184.
- Strickland, J.D.H., Parsons, T.R., 1972. A practical handbook of seawater analysis. Bulletin 167 (Second Edition). Fisheries Research Board of Canada 167, 1–310.
- Sutherland, T.F., Shepherd, P.C.F., Elner, R.W., 2000. Predation on meiofaunal and macrofaunal invertebrates by western sandpipers (*Calidris mauri*): evidence for dual foraging modes. Marine Biology 137(5–6), 983–993.
- Sutherland, T.F., Levings, C.D., Petersen, S.A., Poon, P., Piercey, B., 2007. The use of meiofauna as an indicator of benthic organic enrichment associated with salmonid aquaculture. Marine Pollution Bulletin 54, 1249–1261.
- Swedmark, B., 1964. The interstitial fauna of marine sand. Biological Reviews 39, 1–42.
- ter Braak, C.J.F., 1986. Canonical correspondence analysis: a new eigen vector technique for multivariate direct gradient analysis. Ecology 67, 1167–1179.
- Thiel, V.H., 1966. Quantitative investigations of deep-sea meiofauna. Veroff. Inst. Meeresforsch., Bremerh 11, 131–147.
- Thilagavathi, B., Varadharajan, D., Manoharan, J., Vijayalakshmi, S., Balasubramanian, T., 2012. Taxonomy and distribution of benthic foraminifera from the sediment of Palk Strait, South East coast of India. International Journal of Pharmaceutical & Biological Archive 3, 1129–1136.
- Thistle, D., Sherman, K.M., 1985. The nematode fauna of a deep-sea site exposed to strong near-bottom currents. Deep Sea Research Part A 32(9), 1077–1088.
- Thistle, D., Lambshead, P.J.D., Sherman, K.M., 1995. Nematode tail-shape groups respond to environmental differences in the deep sea. Vie Milieu 45, 107–115.
- Tietjen, J.H., 1969. The ecology of shallow water meiofauna in two New England estuaries. Oecologia 2, 251–291.
- Tietjen, J.H., 1980. Microbial-meiofaunal interrelationships: A review. Microbiology 335-338.

- Torres\_Pratts, H., Schizas, N.V., 2007. Meiofaunal Colonization of Decaying Leaves of the Red Mangrove *Rhizophora mangle*, in Southwestern Puerto Rico, Caribbean Journal of Science 43(1), 127–137.
- UNEP WCMC 1987 (updated May 2011) Sundarbans National Park, West Bengal, India; UNEP World Conservation Monitoring Centre, 11p.
- Vafeiadou, A.M., Chintiroglou, C., Moens, T., 2018a. Effects of an increased temperature regime on the population dynamics and species interactions of marine nematodes. Journal of Experimental Marine Biology and Ecology 502, 142–152.
- Vafeiadou, A.M., Bretaña, B.L.P., Van Colen, C., dos Santos, G.A.P., Moens, T., 2018b. Global warming-induced temperature effects to intertidal tropical and temperate meiobenthic communities. Marine Environmental Research 142, 163–177.
- Vanaverbeke, J., Soetaert, K., Heip, C., Vanreusel, A., 1997. The metazoan meiobenthos along the continental slope of the Goban Spur (NE Atlantic). Journal of Sea Research 38, 93–107.
- Vanaverbeke, J., Steyaert, M., Vanreusel, A., Vincx, M., 2003. Nematode biomass spectra as descriptors of functional changes to human and natural impact. Marine Ecology Progress Series 249, 17–157.
- Vanaverbeke, J., Steyaert, M., Soetaert, K., Rousseau, V., Van Gansbeke, D., Parent, J.Y., Vincx, M., 2004. Changes in structural and functional diversity of nematode communities during a spring phytoplankton bloom in the southern North Sea. Journal of Sea Research 52, 281–292.
- Vanaverbeke, J., Merckx, B., Degraer, S., Vincx, M., 2011. Sediment-related distribution patterns of nematodes and macrofauna: two sides of the benthic coin? Marine Environmental Research 71, 31–40.
- Vanhove, S., Vincx, M., Vangansbeke, D., Gijselinck, W., Schram, D., 1992. The meiobenthos of 5 mangrove vegetation Types in Gazi Bay, Kenya. Hydrobiologia 247, 99–108.

- Vanhove, S., Wittooeck, J., Beghyn, M., Van Gansbeke, D., Van Kenhove, A., Coomans, A., Vincx, M., 1997. Role of meiobenthos in Antartic ecosystems. Belgian Antartic activities. Marine Ecology Progress Series 127, 1–59.
- Vanhove, S., Beghyn, M., Van Gansbeke, D., Bullough, L.W., Vincx, M., 2000. A seasonally varying biotope at Signy Island, Antarctic: implications for meiofaunal structure. Marine Ecology Progress Series 202, 13–25.
- Vanhove, S., Vermeeren, H., Vanreusel, A., 2004. Meiofauna towards the South Sandwich Trench (750–6300 m), focus on nematodes. Deep-Sea Research II 51, 1665–1687.
- Vanreusel, A., 1990. Ecology of the free-living marine nematodes from the Voordelta (Southern Bight of the North Sea). I. Species composition and structure of nematode communities. Cahiers de Biologie Marine 31, 439–462.
- Veiga, P., Besteiro, C., Rubal, M., 2010. Meiofauna communities in exposed sandy beaches on the Galician coast (NW Spain), six months after the Prestige oil spill: the role of polycyclic aromatic hydrocarbons (PAHs). Scientia Marina 74, 385–394.
- Venkataramana, V., Sarma, V.V.S.S., Reddy, A.M., 2017. Impact of river discharge on distribution of zooplankton biomass, community structure and food web dynamics in the Western coastal Bay of Bengal. Regional Studies in Marine Science 16, 267– 278.
- Vincx, M., 1996. Meiofauna in marine and brackish water sediments. In: Hall, G.S., (Eds.), Methods for Examination of Organismal Diversity in Soils and Sediment. CAB International, pp. 187–195.
- Wang, X., Liu, X., Xu, J., 2019. Distribution patterns of meiofauna assemblages and their relationship with environmental factors of deep sea adjacent to the Yap Trench, Western Pacific Ocean. Frontiers in Marine Science 6, 735.
- Warnock, A.M., Hagen, S.C., Passeri, D.L., 2015. Marine Tar Residues: a Review. Water, Air, Soil Pollution 226(3), 68.
- Warwick, R.M., Price, R., 1979. Ecological and metabolic studies on free-living marine nematodes from an estuarine mud-flat. Estuarine, Coastal and Shelf Science 9, 257– 271.

- Warwick, R.M., 1981. The influence of temperature and salinity on energy partitioning in the marine nematode *Diplolaimelloides bruciei*. Oecologia 51, 318–325.
- Warwick, R.M., 1989. Role of meiofauna in the marine ecosystem: Evolutionary considerations. Zoological Journal of the Linnean Society 96, 229–241.
- Warwick, R.M., Platt, H.M., Somerfield, P.J.D., 1998. Free-living marine nematodes, Part III, Monhysterids, Pictorial key to world genera and notes for the identification of British species. In: Kermack, D.M., Barnes, R.S.K., (Eds.), Synopses of the British Fauna (New series). Field Studies Council, Shrewsbury, UK, pp. 1–296.
- Wells, J.B.J., Rao, G.C., 1987. Littoral Harpacticoida (Crustacea: Copepoda) from Andaman and Nicobar Islands. Memoirs of the Zoological Survey of India 16(4), 1– 385.
- Wernberg, T., Russell, B.D., Moore, P.J., Ling, S.D., Smale, D.A., Campbell, A., Coleman, M.A., Steinberg, P.D., Kendrick, G.A., Connell, S.D., 2011. Impacts of climate change in a global hotspot for temperate marine biodiversity and ocean warming. Journal of Experimental Marine Biology and Ecology 400, 7–16.
- Werner, K.M., Perez, L.J., Ghosh, R., Semmelhack, M.F., Bassler, B.L., 2014. *Caenorhabditis elegans* recognizes a bacterial quorum-sensing signal molecule through the AWCON neuron. Journal of Biological Chemistry 289(38), 26566– 26573.
- Wetzel, M.A., Jensen, P., Giere, O., 1995. Oxygen/sulphide regime and nematode fauna associated with *Arenicola marina* burrows: new insights in the thiobios case. Marine Biology 124, 301–312.
- Wetzel, M.A., Weber, A., Giere, O., 2002. Re-colonization of anoxic/sulfidic sediments by marine nematodes after experimental removal of macroalgal cover. Marine Biology 141, 679–689.
- Widdicombe, S., Dashfield, S.L., McNeill, C.L., Needham, H.R., Beesley, A., McEvoy, A., Oxnevad, S., Clarke, K.R., Berge, J.A., 2009. Effects of CO<sub>2</sub> induced seawater acidification on infaunal diversity and sediment nutrient fluxes. Marine Ecology Progress Series 379, 59–75.

- Wieser, W., 1953. Die Beziehung zwischen Mundhöhlengestalt, Ernährungswiese und 1000 Vorkommen bei freilebenden marinen Nematoden. Arkiv för zoology 4, 439–484.
- Wigley, R.L., McIntyre, A.D., 1964. Some quantitative comparisons of offshore meiobenthos and macrobenthos south of Martha's Vineyard. Limnology and Oceanography 9, 485–493.
- Xuan, Q.N., Vanreusel, A., Thanh, N.V., Smol, N., 2007. Biodiversity of meiofauna in the intertidal Khe Nhan mud at, Can Gio mangrove forest, Vietnam with special emphasis on free living nematodes. Ocean Science Journal 42(3), 135–152.
- Zeppilli, D., Sarrazin, J., Leduc, D., Arbizu, P.M., Fontaneto, D., Fontanier, C., Gooday, A.J., Kristensen, R.M., Ivanenko, V.N., Sorensen, M.V., Vanreusel, A., Thébault, J., Mea, M., Allio, N., Andro, T., Arvigo, A., Castrec, J., Danielo, M., Foulon, V., Fumeron, R., Hermabessiere, L., Hulot, V., James, T., Langonne-Augen, R., Le Bot, T., Long, M., Mahabror, D., Morel, Q., Pantalos, M., Pouplard, E., Raimondeau, L., Rio-Cabello, A., Seite, S., Traisnel, G., Urvoy, K., Van Der Stegen, T., Weyand, M., Fernandes, D., 2015. Is the meiofauna a good indicator for climate change and anthropogenic impacts? Marine Biodiversity 45, 505–535.
- Zeppilli, D., Leduc, D., 2018. Biodiversity and ecology of meiofauna in extreme and changing environments. Marine Biodiversity 48, 1–4.
- Zhou, H., 2001. Effects of leaf litter addition on meiofaunal colonization of azoic sediments in a subtropical mangrove in Hong Kong. Journal of Experimental Marine Biology and Ecology 256, 99–121.