

## Table of contents:

Titles

Table of content

Table of Abbreviations

Tables of Figures

### **Introduction chapter:**

A. Cancer overview.....	16-17
B. Types of Cancer.....	17-18
C. Types of Leukemia.....	18-20
D. Chronic Myeloid Leukemia.....	20-22
E. Therapy for CML.....	22-24
F. Inhibitor of apoptotic protein and its role.....	24-27
G. Targeting the IAPs for cancer therapy.....	27-31
H. Immunotherapy.....	31
I. TRAIL therapy.....	32
J. Reactive oxygen species.....	33
K. Redox status in haematological malignancies.....	32-33
L. Mechanism of ROS production in cells.....	34-35
M. Redox status: a biological basis for therapeutic selectivity.....	35
N. ROS mediated Signalling Pathway.....	35-39
O. Mechanism of chemoresistance different from chemosensitive.....	39-44

### **Chapter 1: Identification of Inhibitor of apoptotic protein as a target of induction of apoptosis in imatinib resistant CML cells**

Introduction.....	46-47
Material and Methods.....	47-49
Results	
A. Imatinib-resistant CML cell lines K562(R) show resistance against TRAIL-induced apoptosis.....	49-50
B. Hydroxychavicol sensitizes imatinib-resistant CML cell lines to TRAIL-induced apoptosis.....	51-52
C. Involvement of TRAIL mediated extrinsic apoptotic pathway in Hydroxychavicol and TRAIL combinatory treatment in K562(R) cells.....	53
D. Hydroxychavicol did not alter DR4 and DR5 expression and DCR1 and DCR2 expression at RNA level and Protein level.....	53-56
E. Hydroxychavicol decreased XIAP and FLIP at the protein level.....	56-57
F. Proteosomal mediated degradation pathway is involved in Hydroxychavicol mediated XIAP and FLIP degradation.....	58
G. Knockdown of XIAP and FLIP involved in TRAIL mediated apoptosis.....	59-60

H. ROS played a major role in Hydroxychavicol -mediated TRAIL sensitization...	60-61
I. Hydroxychavicol only increases intracellular ROS level in Imatinib resistant K562 cells.....	61-63
J. Hydroxychavicol downregulates XIAP and FLIP in ROS dependant.....	63-64
K. ROS is involved in Hydroxychavicol mediated TRAIL induced apoptosis.....	65-66
Discussion.....	66-68

**Chapter 2:** To identify the modifications of IAPs and signalling mechanisms involved in the modification

Introduction.....	70-71
Materials and Methods.....	71-74
Results	
A. Hydrogen Peroxide Down-regulates FLIP and XIAP.....	74-75
B. H <sub>2</sub> O <sub>2</sub> does not cause cell cytotoxicity on K562 (R) cells.....	75-76
C. H <sub>2</sub> O <sub>2</sub> did not downregulate XIAP and FLIP at RNA level.....	76-77
D. H <sub>2</sub> O <sub>2</sub> solely involved in XIAP and FLIP downregulation.....	77-78
E. XIAP and cFLIP degradation is proteasomal dependent.....	78-80
F. XIAP and cFLIP degradation is not lysosomal dependent.....	80-81
G. Activation of ERK and JNK plays role in XIAP, FLIP downregulation respectively.....	81-83
H. H <sub>2</sub> O <sub>2</sub> increases JNK and ERK activation by phosphorylation dependant manner.....	83-85
I. JNK, ERK played differential role in downregulation in cellular level of FLIP, XIAP respectively.....	85-88
J. ERK mediated PI3K and Akt dephosphorylation involved in XIAP ubiquitination and degradation.....	89-92
K. Role of ERK in PI3K and Akt dephosphorylation.....	92-93
L. JNK not involved in Akt and PI3K dephosphorylation.....	94
M. ITCH is responsible for FLIP ubiquitination and degradation.....	96-97
N. JNK is involved in ITCH mediated FLIP ubiquitination and downregulation.....	97-98
O. Discussion.....	101

**Chapter 3:** To check the key findings in imatinib resistant cells from objective I and objective II in imatinib sensitive cells

Introduction.....	103
Material and Method.....	103-105
Results	
A. CML cell lines show resistance against TRAIL-induced apoptosis .....	105-106

B. Hydroxychavicol sensitizes CML cell lines to TRAIL-induced apoptosis.....	<b>106-107</b>
C. Hydroxychavicol-mediated TRAIL sensitization of K562(S) cell is ROS dependent: .....	<b>108-109</b>
D. Hydroxychavicol decreased XIAP and FLIP at the protein level.....	<b>109-110</b>
E. Downregulation of antiapoptotic proteins XIAP and FLIP happens in a ROS-dependent manner.....	<b>110-112</b>
F. Externally treated H <sub>2</sub> O <sub>2</sub> do not alter XIAP and FLIP at mRNA level in imatinib sensitive K562 cells .....	<b>113</b>
G. H <sub>2</sub> O <sub>2</sub> downregulated FLIP and XIAP via Lysosomal dependant pathway not proteosomal pathway.....	<b>114-115</b>
H. H <sub>2</sub> O <sub>2</sub> does not cause significant apoptosis as compare to control in imatinib sensitive K562 cells: .....	<b>116</b>
I. MAPK signaling pathway regulates Hydroxychavicol mediated TRAIL induced apoptosis.....	<b>117</b>
Discussion.....	<b>118</b>

#### **Chapter 4**

Overall Discussion and conclusion.....	<b>120-127</b>
Future Direction and Therapeutical impact of study.....	<b>128</b>